



United States Department of Agriculture

Smoke Management Photographic Guide: A Visual Aid for Communicating Impacts

Joshua C. Hyde, Jarod Blades, Troy Hall, Roger D. Ottmar, and Alistair Smith



Forest Service

Pacific Northwest Research Station

General Technical Report
PNW-GTR-925

May
2016

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.



Authors

Joshua C. Hyde is a program coordinator, University of Idaho, College of Natural Resources, 400 North 34th Street, Suite 201, Seattle, WA 98103; **Jarod Blades** is an assistant professor, University of Wisconsin, Plant and Earth Sciences, 410 South Third Street, River Falls, WI 54022; **Troy Hall** is the head of the Department of Conservation Social Sciences, Oregon State University, College of Forestry, 321 Richardson Hall, Corvallis, OR 97331; **Roger D. Ottmar** is a research forester, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 400 North 34th Street, Suite 201, Seattle, WA 98103; **Alistair Smith** is an associate professor, University of Idaho, College of Natural Resources, 975 West Sixth Street, Moscow, ID 83844.

Published in cooperation with: the University of Idaho—College of Natural Resources and the Joint Fire Science Program.

Abstract

Hyde, Joshua C.; Blades, Jarod; Hall, Troy; Ottmar, Roger D.; Smith, Alistair.

2016. Smoke management photographic guide: a visual aid for communicating impacts. Gen. Tech. Rep. PNW-GTR-925. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 59 p.

Communicating emissions impacts to the public can sometimes be difficult because quantitatively conveying smoke concentrations is complicated. Regulators and land managers often refer to particulate-matter concentrations in micrograms per cubic meter, but this may not be intuitive or meaningful to everyone. The primary purpose of this guide is to serve as a tool for communicating potential particulate matter (PM_{2.5}) levels during wildfire events using visual representation. Examples of visibility impairment under various levels of smoke concentration and humidity have been modeled using the WinHaze program.

Keywords: Air quality, regional haze, smoke management.

Contents

- 1 **Introduction**
- 1 **Methods**
- 3 **Using This Guide**
- 4 **Limitations**
- 5 **English Equivalent**s
- 5 **Acknowledgments**
- 5 **Literature Cited**
- 8 **Appendix**
- 8 **Northern Region (National Forest System—Region 1):
Selway-Bitterroot Wilderness and Glacier National Park (Montana)**
- 14 **Intermountain Region (National Forest System—Region 2):
Rocky Mountain National Park (Colorado)**
- 18 **Southwestern Region (National Forest System—Region 3):
Grand Canyon National Park (Arizona)**
- 24 **Rocky Mountain Region (National Forest System—Region 4):
Canyonlands National Park (Utah)**
- 28 **Rocky Mountain Region (National Forest System—Region 4):
Great Basin National Park (Nevada)**
- 32 **Pacific Southwest Region (National Forest System—Region 5):
Yosemite National Park (California)**
- 36 **Pacific Northwest Region (National Forest System—Region 6):
Columbia River Gorge National Scenic Area (Oregon)**
- 40 **Pacific Northwest Region (National Forest System—Region 6):
Snoqualmie Pass (Washington)**

- 44 **Southern Region (National Forest System—Region 8):
Great Smoky Mountains National Park (Tennessee)**
- 48 **Southern Region (National Forest System—Region 8):
Mammoth Cave National Park (Kentucky)**
- 52 **Southern Region (National Forest System—Region 8):
Big Bend National Park (Texas)**
- 56 **Eastern Region (National Forest System—Region 9):
Acadia National Park (Maine)**

Introduction

An important but difficult part of determining the impacts of emissions from wildland fire on air quality is the preparation of a quantitative smoke-concentration assessment. Although smoke is composed of a variety of chemical compounds and other components, regulators and land managers often focus on particulate matter (PM) owing to its effects on human health and visibility degradation. Particulate matter in smoke generally ranges from 0.1 to 100 micrometers (μm) in diameter (Hardy et al. 2001). Diameters $\leq 10 \mu\text{m}$ (PM_{10}) and $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) are the most common size classes used in air quality measurement and monitoring. Particulate matter concentration is measured in units of micrograms per cubic meter ($\mu\text{g}/\text{m}^3$).

Visibility is affected by several factors, including the composition and concentration of smoke from fires. Particulate matter emitted from fires can contain ammonium nitrate, ammonium sulfate, and light-absorbing carbon (Malm et al. 1994). The effect these compounds have on visibility can be magnified by increases in relative humidity, which causes more water vapor to adhere to particles thereby altering the way they absorb and reflect light (Malm et al. 2003). In addition, background levels of visibility vary geographically (Hand et al. 2014, Malm et al. 1994).

This guide illustrates the effects of wildland fire smoke on visibility for the continental United States (fig. 1). It was developed with images from locations in national parks and other scenic areas to assess visibility impairment associated with elevated $\text{PM}_{2.5}$ concentration through visual representation. Images presented in this guide were generated using WinHaze (Air Resource Specialists, Inc. 2013), a software tool developed to visualize the impacts of pollution on visibility.¹ Because of the complex relationships that influence observed visual range approximations—contrasts perceived by the naked eye, the effects of these contrasts on perception of visual range, and the effects of both of these factors on particle-concentration estimates—they should be used as general indicators, not precise measurements.

Methods

This reference guide was generated using WinHaze imaging software version 2.9.9.1. (Air Resource Specialists, Inc. 2013) for the purpose of representing visual impacts from smoke in numerous U.S. locations. To determine the reduction in visibility, WinHaze incorporates several years of particulate monitoring data and images from national parks and wilderness areas into a beta extinction equation

¹ The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

**This guide will help
air quality regulators
and land managers
communicate $\text{PM}_{2.5}$
concentrations during
wildland fires.**

- Northern (Region 1): Selway-Bitterroot Wilderness (south), Glacier National Park (north)
- Intermountain (Region 2): Rocky Mountain National Park
- Southwestern (Region 3): Grand Canyon National Park (both)
- Rocky Mountain (Region 4): Canyonlands National Park (east), Great Basin National Park (west)
- Pacific Southwest (Region 5): Yosemite National Park
- Pacific Northwest (Region 6): Columbia River Gorge National Scenic Area (south), Snoqualmie Pass (north)
- Southern (Region 8): Great Smoky Mountains National Park (east), Mammoth Cave National Park (center), Big Bend National Park (west)
- Eastern (Region 9): Acadia National Park




Figure 1—Locations of sites where estimations of decreased visibility were converted into photo images (grey dots). The sites selected represent typical conditions in each of the National Forest System regions of the contiguous U.S. states.

(Hand and Malm 2006); the equation, images, and data are products of the Inter-agency Monitoring of Protected Visual Environments (IMPROVE) program, which has placed stationary cameras and air quality monitoring equipment at several national parks (<http://vista.cira.colostate.edu/improve/Default.htm>. [15 June 2015]).

WinHaze allows for consistent visual representation of air quality under varying levels of humidity, background pollution, and $PM_{2.5}$ concentration. Each example in this guide presents a simulated baseline image that represents the visual range under average PM concentrations juxtaposed with several images of simulated visual impairment. Visual impairment from smoke was simulated first by establishing constant values for the $PM_{2.5}$ constituents that would be unlikely to change with the introduction of smoke from fires; values for ammonium sulfate, ammonium nitrate, and fine soil were calculated by averaging the 20 percent worst visibility days. Organic carbon and black carbon—which account for nearly 75 percent of the emissions from forest fires (Andreae and Merlet 2001)—were then increased to reflect increasing concentrations of smoke using an organic-carbon to black-carbon ratio of 15.4:1 (Andreae and Merlet 2001). The coarse particulate inputs used to simulate each image were chosen based on the larger of two values: either the average value of the 20 percent worst monitored days, or 10 percent of the $PM_{2.5}$ (ammonium sulfate, ammonium nitrate, fine soil, organic carbon, and black carbon) concentration (Ward and Hardy 1991).

The version of WinHaze used for this work includes the first version of the IMPROVE beta extinction equation, as described in Hand and Malm (2006). To

improve the accuracy of the simulations by accounting for hygroscopicity (Malm et al. 2005)—the ability of a substance to attract and hold water molecules from the surrounding environment—a correction factor was applied to the organic carbon values before each image and visual range determination was generated; accordingly, light scattering (total beta extinction) of organic carbon increased linearly by a factor of 1.2 at 80-percent relative humidity compared to no relative humidity. Each image includes prominent landmarks with which to judge visual range. The distance between the camera locations and each landmark was measured with Google Earth and verified using location information from Air Resource Specialists, Inc.

Relative humidity affects visibility and changes throughout the day and from one season to the next; therefore, a range of values was chosen to represent morning and afternoon monthly averages most likely to occur during the wildland fire season (May to September) in all national park and wilderness-area locations (EPA 2014). Because these data were unavailable for the two Pacific Northwest (Region 6) locations (fig. 1), meteorological station data were chosen from a location as geographically close to the available site as possible (NOAA 2014).

Lipsett et al. (2012) and the Environmental Protection Agency (EPA 2013) define five levels for air quality: good, moderate, unhealthy for sensitive groups, unhealthy, and very unhealthy (table 1). These levels correspond to thresholds for action by public health officials: good requires no action, moderate suggests sensitive populations reduce prolonged or heavy exertion, unhealthy for sensitive groups requires warnings or alerts to those with heart or lung conditions or other pertinent health issues, unhealthy requires that all people should be notified, regardless of health status, and very unhealthy recommends everyone should avoid physical activity outdoors (Lipsett et al. 2012). The $PM_{2.5}$ levels that were chosen for display in this guide were good ($<38 \mu\text{g}/\text{m}^3$), unhealthy for sensitive groups (89 to $138 \mu\text{g}/\text{m}^3$), and unhealthy (139 to $351 \mu\text{g}/\text{m}^3$) for a short period of time (up to 3 hours) because they are sufficiently different as to be easily discernable to the naked eye. The mid-point of each range was chosen to represent each health level: $19 \mu\text{g}/\text{m}^3$ for good, $114 \mu\text{g}/\text{m}^3$ for unhealthy for sensitive groups, and $245 \mu\text{g}/\text{m}^3$ for unhealthy.

Using This Guide

Each set of images in this guide is preceded by a description of the air quality data for the site depicted, including the date range and number of sampling days of PM data used by WinHaze, the source for the relative humidity data, a table listing the constituents of smoke (both $PM_{2.5}$ and PM_{10}) represented in the images, and a table

Each set of images includes tables documenting the constituents of particulate matter (PM) at different concentration levels, and visual range at different relative humidity and PM concentrations levels.

Public health officials may recommend different actions based on the concentration of smoke in the area.

Table 1—Images and visual range estimates representing the particulate matter (PM) concentration mid-points of the good (19 $\mu\text{g}/\text{m}^3$), unhealthy for sensitive groups (114 $\mu\text{g}/\text{m}^3$), and unhealthy (245 $\mu\text{g}/\text{m}^3$) categories were chosen for display in this guide

Air quality	PM ₁₀ or PM _{2.5} concentration ($\mu\text{g}/\text{m}^3$) ^{a,b}	Actions required to protect health
Good	0 to 38	- No action needed.
Moderate	39 to 88	- Unusually sensitive people should consider reducing prolonged or heavy exertion.
Unhealthy for sensitive groups	89 to 138	- People with heart or lung disease, children, and older adults should reduce prolonged or heavy outdoor exertion. - Everyone else should limit prolonged or heavy exertion.
Unhealthy	139 to 351	- People with heart or lung disease, children, and older adults should avoid all physical activity outdoors. - Everyone else should avoid prolonged or heavy exertion.
Very unhealthy	>351	- People with heart or lung disease, children, and older adults should remain indoors and keep activity levels low. - Everyone else should avoid all physical activity outdoors.

^a PM₁₀ and PM_{2.5} are composed of particles that are ≤ 10 μm and ≤ 2.5 μm in diameter, respectively.

^b Concentrations are 1- to 3-hour averages.

Source: Adapted from Lipsett et al. (2012) and EPA (2013).

listing all of the visible range distances for each PM_{2.5} concentration and relative humidity level.

To use this guide for visualizing PM_{2.5} concentration, select the region and location that best matches the terrain and humidity conditions of the location you are assessing and compare your line-of-sight with landmarks located at distances that correspond to those shown in the photographs. Unless no distinction could be made between photographs, as sometimes occurs at the higher PM_{2.5} concentration levels, images are included that represent baseline (smoke free), good (19 $\mu\text{g}/\text{m}^3$), unhealthy for sensitive groups (114 $\mu\text{g}/\text{m}^3$), and unhealthy (245 $\mu\text{g}/\text{m}^3$) conditions for each location.

Limitations

Visual range is simulated based on analyses of both air quality data and the constituents of wildland fire smoke. Images included in this guide were generated

independently of one factor (sun angle) that has an effect on visibility (Malm and Schichtel 2013, Middleton 1968). Also, the PM_{2.5} concentration levels for good, unhealthy for sensitive groups, and unhealthy conditions shown in the photographs are not instantaneous “snapshots,” but are based on average levels over a period of 1 to 3 hours. This is important because visual ranges can change relatively rapidly.

English Equivalent

When you know:	Multiply by:	To find:
Micrometers (μm)	0.039	Mils
Micrograms (μg)	0.00002	Grains
Kilometers (km)	0.62	Miles

Acknowledgments

We acknowledge funding from the Joint Fire Science Program under Project JFSP 10-1-03-2. The authors would also like to thank members of the National Wildfire Coordinating Group Smoke Committee for their comments and feedback during the development of this guide. Mark Fitch, William Malm, Susan O’Neill, and Janice Peterson provided helpful reviews. Robert Vihnanek, Carol Whitlock, and Clint Wright provided production and editorial assistance.

Literature Cited

Air Resource Specialists, Inc. 2013. WinHaze. Version 2.9.9.1. <http://www.air-resource.com/resources/downloads.html>. (20 June 2015).

Andraea, M.O.; Merlet, P. 2001. Emission of trace gases and aerosols from biomass burning. *Global Biogeochemical Cycles*. 15(4): 955–966.

Hand, J.L.; Malm, W.C. 2006. Review of the IMPROVE equation for estimating ambient light extinction coefficients—final report. http://vista.cira.colostate.edu/improve/publications/graylit/016_IMPROVEeqReview/IMPROVEeqReview.htm. (17 June 2015).

Hand, J.L.; Schichtel, B.A.; Malm, W.C. [et al.]. 2014. Widespread reductions in haze across the United States from the early 1990s through 2011. *Atmospheric Environment*. 94: 671–679.

Hardy, C.C.; Ottmar, R.D.; Peterson, J.L.; Core, J.E.; Seamon, P., eds. 2001. Smoke management guide for prescribed and wildland fire: 2001 edition. NFES 1279. Boise, ID: National Interagency Fire Center, National Wildfire Coordinating Group, Fire Use Working Team. 226 p.

- Lipsett, M.; Materna, B.; Stone, S.L. [et al.]. 2012.** Wildfire smoke: a guide for public health officials. Rev. U.S. Environmental Protection Agency. 53 p. http://oehha.ca.gov/air/risk_assess/wildfirev8.pdf. (17 June 2015).
- Malm, W.C.; Day, D.E.; Kreidenweis, S.M.; Collett, J.L., Jr.; Lee, T. 2003.** Humidity-dependent optical properties of fine particles during the Big Bend Regional Aerosol and Visibility Observational Study. *Journal of Geophysical Research*. 108(D9): 4279.
- Malm, W.C.; Day, D.E.; Kreidenweis, S.M. [et al.]. 2005.** Hygroscopic properties of an organic-laden aerosol. *Atmospheric Environment*. 39: 4969–4982.
- Malm, W.C.; Schichtel, B.A. 2013.** Uncertainty associated with estimating a short-term (1–3 hr) particulate matter concentration from a human-sighted visual range. Joint Fire Science Program Project Report. 105 p. <http://digitalcommons.unl.edu/jfस्पresearch/5/>. (17 June 2015).
- Malm, W.C.; Sisler, J.F.; Huffman, D.; Eldred, R.A.; Cahill, T.A. 1994.** Spatial and seasonal trends in particle concentration and optical extinction in the United States. *Journal of Geophysical Research*. 99(D1): 1347–1370.
- Middleton, W.E.K. 1968.** Vision through the atmosphere; corrected edition. Toronto, Canada: University of Toronto Press. 250 p.
- National Oceanic and Atmospheric Administration [NOAA]. 2014.** Quality controlled local climatological data. Version 2.5.5. <http://cdo.ncdc.noaa.gov/qclcd/QCLCD?prior=N>. (20 June 2015).
- U.S. Environmental Protection Agency [EPA]. 2013.** Air quality guide for particle pollution. 2 p. http://www.epa.gov/airnow/air-quality-guide_pm_2013.pdf. (17 June 2015).
- U.S. Environmental Protection Agency [EPA]. 2014.** Clean air status and trends network (CASTNET). <http://epa.gov/castnet/javaweb/index.html>. (21 January 2015).
- Ward, D.E.; Hardy, C.C. 1991.** Smoke emissions from wildland fires. *Environment International*. 17(2–3): 117–134.

Appendix

Northern Region (National Forest System—Region 1): Selway-Bitterroot Wilderness and Glacier National Park (Montana)

Particulate data from 1,037 days of sampling (March 1988 to May 1999) at Glacier National Park were chosen to represent baseline and elevated regional air quality concentrations (table 2). Table 3 shows the simulated visual range at different levels of $PM_{2.5}$ concentration (<5, 19, 114, and 245 $\mu\text{g}/\text{m}^3$) and relative humidity for both the Selway-Bitterroot Wilderness and Glacier National Park. The simulated images show a baseline representing an area free of smoke-impaired visibility (<5 $\mu\text{g}/\text{m}^3$ fine and coarse particulates) and two or three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Glacier National Park (EPA 2014).

Table 2—Constituents of particulate matter ($PM_{2.5}$ and PM_{10}) at baseline (<5 $\mu\text{g}/\text{m}^3$) and elevated (19, 114, and 245 $\mu\text{g}/\text{m}^3$) levels in the Selway-Bitterroot Wilderness and Glacier National Park, Montana

Particulate matter constituents	Particulate matter concentration			
	<5 (Baseline)	19	114	245
		$\mu\text{g}/\text{m}^3$		
Ammonium sulfate	0.96	1.29	1.29	1.29
Ammonium nitrate	0.30	0.61	0.61	0.61
Organic carbon	2.67	14.95	104.14	227.13
Black carbon	0.43	0.97	6.78	14.79
Fine soil	0.58	1.19	1.19	1.19
Coarse mass	6.12	10.21	11.40	24.50



Note: PM_{10} and $PM_{2.5}$ are composed of particles that are $\leq 10 \mu\text{m}$ and $\leq 2.5 \mu\text{m}$ in diameter, respectively.



Table 3—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in the Selway-Bitterroot Wilderness and Glacier National Park, Montana

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	Percent	Miles	Kilometers
<5 (Baseline)	40	72.7	117.0
	60	23.1	37.2
19 (Good)	40	24.6	39.6
	60	23.1	37.2
	80	21.3	34.3
	90	19.9	32.1
	90	19.9	32.1
114 (Unhealthy for Sensitive Groups)	40	4.4	7.1
	60	4.2	6.8
	80	4.0	6.5
	90	3.9	6.3
	90	3.9	6.3
245 (Unhealthy)	40	2.1	3.3
	50 to 60	2.0	3.2
	70 to 90	1.9	3.1



^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.


Selway-Bitterroot Wilderness (Montana)





Glacier National Park (Montana)





Intermountain Region (National Forest System—Region 2): Rocky Mountain National Park (Colorado)

Particulate data from 794 days of sampling (September 1990 to May 1999) at Rocky Mountain National Park were chosen to represent baseline and elevated regional air quality concentrations (table 4). Table 5 shows the simulated visual range at different levels of $PM_{2.5}$ concentration (<5, 19, 114, and 245 $\mu\text{g}/\text{m}^3$) and relative humidity for Rocky Mountain National Park. The simulated images show a baseline representing an area free of smoke-impaired visibility (<5 $\mu\text{g}/\text{m}^3$ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Rocky Mountain National Park (EPA 2014).

Table 4—Constituents of particulate matter ($PM_{2.5}$ and PM_{10}) at Baseline (<5 $\mu\text{g}/\text{m}^3$) and elevated (19, 114, and 245 $\mu\text{g}/\text{m}^3$) levels in Rocky Mountain National Park, Colorado

Particulate matter constituents	Particulate matter concentration			
	<5 (baseline)	19	114	245
		$\mu\text{g}/\text{m}^3$		
Ammonium sulfate	0.93	1.49	1.49	1.49
Ammonium nitrate	0.29	0.50	0.50	0.50
Organic carbon	1.00	14.77	103.96	226.95
Black carbon	0.17	0.96	6.77	14.78
Fine soil	0.63	1.28	1.28	1.28
Coarse mass	3.96	5.88	11.40	24.50



Note: PM_{10} and $PM_{2.5}$ are composed of particles that are $\leq 10 \mu\text{m}$ and $\leq 2.5 \mu\text{m}$ in diameter, respectively.



Table 5—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Rocky Mountain National Park, Colorado

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	<i>Percent</i>	<i>Miles</i>	<i>Kilometers</i>
<5 (Baseline)	40	107.5	173.0
19 (Good)	40	25.4	40.9
	50	24.6	39.6
	60	23.8	38.3
114 (Unhealthy for Sensitive Groups)	40	4.4	7.1
	50	4.3	7.0
	60	4.2	6.8
245 (Unhealthy)	40	2.1	3.3
	50 to 60	2.0	3.2

^aPM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Rocky Mountain National Park (Colorado)





Southwestern Region (National Forest System—Region 3): Grand Canyon National Park (Arizona)

Particulate data from 857 days (March 1988 to August 1998) at Grand Canyon National Park were chosen to represent baseline and elevated regional air quality concentrations (table 6). Table 7 shows the simulated visual range at different levels of PM_{2.5} concentration (<3, 19, 114, and 245 µg/m³) and relative humidity for two viewing locations at Grand Canyon National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<3 µg/m³ fine and coarse particulates) and two or three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Grand Canyon National Park (EPA 2014).

Table 6—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<3 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Grand Canyon National Park, Arizona

Particulate matter constituents	Particulate matter concentration			
	<3 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	1.01	1.59	1.59	1.59
Ammonium nitrate	0.20	0.31	0.31	0.31
Organic carbon	0.80	14.90	104.09	227.08
Black carbon	0.18	0.97	6.78	14.79
Fine soil	0.61	1.23	1.23	1.23
Coarse mass	4.99	7.16	11.40	24.50



Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.


Table 7—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Grand Canyon National Park, Arizona

PM _{2.5} concentration ^a	Relative humidity		Visual range	
	Percent	Miles	Kilometers	
<3 (Baseline)	10	109.4	176.0	
19 (Good)	10	26.4	42.5	
	20	26.0	41.8	
	30	25.6	41.2	
	40	25.1	40.4	
	50	24.4	39.2	
	60	23.5	37.9	
114 (Unhealthy for Sensitive Groups)	10	4.7	7.6	
	20	4.6	7.4	
	30	4.5	7.3	
	40	4.4	7.1	
	50	4.3	7.0	
	60	4.2	6.8	
245 (Unhealthy)	10	2.2	3.5	
	20 to 40	2.1	3.4	
	50	2.0	3.2	
	60	1.9	3.1	



^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.



Grand Canyon National Park (Arizona)—Bluff View





Grand Canyon National Park (Arizona)—Valley View





Rocky Mountain Region (National Forest System—Region 4): Canyonlands National Park (Utah)

Particulate data from 964 days (March 1988 to May 1999) at Canyonlands National Park were chosen to represent baseline and elevated regional air quality concentrations (table 8). Table 9 shows the simulated visual range at different levels of PM_{2.5} concentration (<3, 19, 114, and 245 µg/m³) and relative humidity for Canyonlands National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<3 µg/m³ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Canyonlands National Park (EPA 2014).

Table 8—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<3 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Canyonlands National Park, Utah

Particulate matter constituents	Particulate matter concentration			
	<3 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	1.08	1.54	1.54	1.54
Ammonium nitrate	0.23	0.37	0.37	0.37
Organic carbon	0.82	14.77	103.96	226.95
Black carbon	0.16	0.96	6.77	14.78
Fine soil	0.69	1.36	1.36	1.36
Coarse mass	5.60	8.43	11.40	24.50



Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.

Table 9—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Canyonlands National Park, Utah

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	<i>Percent</i>	<i>Miles</i>	<i>Kilometers</i>
<3 (Baseline)	10	106.3	171.0
	20	26.3	42.4
19 (Good)	20	26.0	41.8
	30	25.5	41.1
	40	25.0	40.3
	10	4.7	7.6
114 (Unhealthy for Sensitive Groups)	20	4.6	7.4
	30	4.5	7.3
	40	4.4	7.1
	10	2.2	3.5
245 (Unhealthy)	20 to 40	2.1	3.4

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Canyonlands National Park (Utah)





Rocky Mountain Region (National Forest System—Region 4): Great Basin National Park (Nevada)

Particulate data from 681 days of sampling (May 1992 to May 1999) at Great Basin National Park were chosen to represent baseline and elevated regional air quality concentrations (table 10). Table 11 shows the simulated visual range at different levels of PM_{2.5} concentration (<3, 19, 114, and 245 µg/m³) and relative humidity for Great Basin National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<3 µg/m³ fine and coarse particulates) and two levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Great Basin National Park (EPA 2014).

Table 10—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<3 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Great Basin National Park, Nevada

Particulate matter constituents	Particulate matter concentration			
	<3 (Baseline)	19	114	245
	<i>µg/m³</i>			
Ammonium sulfate	0.68	1.13	1.13	1.13
Ammonium nitrate	0.16	0.31	0.31	0.31
Organic carbon	0.98	15.18	104.37	227.37
Black carbon	0.19	0.99	6.80	14.81
Fine soil	0.60	1.39	1.39	1.39
Coarse mass	3.73	5.50	11.40	24.50



Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.


Table 11—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Great Basin National Park, Nevada

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	Percent	Miles	Kilometers
<3 (Baseline)	20	115.0	185.0
	30	26.2	42.2
19 (Good)	30	25.8	41.6
	40	25.4	40.8
	50	24.7	39.7
	50	24.7	39.7
114 (Unhealthy for Sensitive Groups)	20	4.6	7.4
	30	4.5	7.3
	40	4.4	7.1
	50	4.3	7.0
245	20 to 40	2.1	3.4
	50	2.0	3.3

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Great Basin National Park (Nevada)





Pacific Southwest Region (National Forest System—Region 5): Yosemite National Park (California)

Particulate data from 951 days (March 1988 to May 1999) at Yosemite National Park were chosen to represent baseline and elevated regional air quality concentrations (table 12). Table 13 shows the simulated visual range at different levels of $PM_{2.5}$ concentration (<5, 19, 114, and 245 $\mu\text{g}/\text{m}^3$) and relative humidity for Yosemite National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<5 $\mu\text{g}/\text{m}^3$ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Yosemite National Park (EPA 2014).

Table 12—Constituents of particulate matter ($PM_{2.5}$ and PM_{10}) at baseline (<5 $\mu\text{g}/\text{m}^3$) and elevated (19, 114, and 245 $\mu\text{g}/\text{m}^3$) levels in Yosemite National Park, California

Particulate matter constituents	Particulate matter concentration			
	<5 (Baseline)	19	114	245
		$\mu\text{g}/\text{m}^3$		
Ammonium sulfate	0.99	1.90	1.90	1.90
Ammonium nitrate	0.47	0.94	0.94	0.94
Organic carbon	1.94	14.20	103.39	226.38
Black carbon	0.27	0.92	6.73	14.74
Fine soil	0.56	1.04	1.04	1.04
Coarse mass	4.78	7.64	11.40	24.50



Note: PM_{10} and $PM_{2.5}$ are composed of particles that are $\leq 10 \mu\text{m}$ and $\leq 2.5 \mu\text{m}$ in diameter, respectively.



Table 13—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Yosemite National Park, California

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	<i>Percent</i>	<i>Miles</i>	<i>Kilometers</i>
<5 (Baseline)	20	85.7	138.0
	50	24.4	39.2
19 (Good)	20	26.2	42.1
	30	25.8	41.5
	40	25.2	40.6
	50	24.4	39.2
	50	24.4	39.2
114 (Unhealthy for Sensitive Groups)	20	4.6	7.4
	30	4.5	7.3
	40	4.4	7.1
	50	4.3	7.0
	50	4.3	7.0
245 (Unhealthy)	20	2.1	3.4
	50	2.0	3.2

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Yosemite National Park (California)





Pacific Northwest Region (National Forest System—Region 6): Columbia River Gorge National Scenic Area (Oregon)

Particulate data from 551 days of sampling (July 1993 to May 1999) in the Columbia River Gorge National Scenic Area were chosen to represent baseline and elevated regional air quality concentrations (table 14). Table 15 shows the simulated visual range at different levels of PM_{2.5} concentration (<6, 19, 114, and 245 µg/m³) and relative humidity for the Columbia River Gorge National Scenic Area. The simulated images show a baseline representing an area free from smoke-impaired visibility (<6 µg/m³ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Portland, Oregon (NOAA 2014).

Table 14—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<6 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in the Columbia River Gorge National Scenic Area, Oregon

Particulate matter constituents	Particulate matter concentration			
	<6 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	1.48	2.56	2.56	2.56
Ammonium nitrate	0.77	1.78	1.78	1.78
Organic carbon	2.32	12.56	101.75	224.75
Black carbon	0.47	0.82	6.63	14.63
Fine soil	0.66	1.28	1.28	1.28
Coarse mass	7.90	11.88	11.88	24.50


Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.



Table 15—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in the Columbia River Gorge National Scenic Area, Oregon

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	<i>Percent</i>	<i>Miles</i>	<i>Kilometers</i>
<6 (Baseline)	40	66.5	107.0
	50	66.5	107.0
19 (Good)	40	25.4	40.8
	50	24.2	39.0
	60	22.9	36.9
	70	21.5	34.6
	80	19.9	32.0
114 (Unhealthy for Sensitive Groups)	40	4.4	7.2
	50	4.3	7.0
	60	4.2	6.8
	70	4.1	6.6
	80	4.0	6.4
245 (Unhealthy)	40	2.1	3.3
	50 to 60	2.0	3.2
	70 to 80	1.9	3.1

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Columbia River Gorge National Scenic Area (Oregon)





Pacific Northwest Region (National Forest System—Region 6): Snoqualmie Pass (Washington)

Particulate data from 353 days of sampling (December 1993 to May 1999) at Snoqualmie Pass were chosen to represent baseline and elevated regional air quality concentrations (table 16). Table 17 shows the simulated visual range at different levels of PM_{2.5} concentration (<4, 19, 114, and 245 µg/m³) and relative humidity for Snoqualmie Pass. The simulated images show a baseline representing an area free from smoke-impaired visibility (<4 µg/m³ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Seattle-Tacoma International Airport, Washington (NOAA 2014).

Table 16—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<4 µg/m³) and elevated (19, 114, and 245 µg/m³) levels at Snoqualmie Pass, Washington

Particulate matter constituents	Particulate matter concentration			
	<4 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	0.98	1.84	1.84	1.84
Ammonium nitrate	0.35	0.59	0.59	0.59
Organic carbon	1.28	14.80	103.99	226.98
Black carbon	0.31	0.96	6.77	14.78
Fine soil	0.29	0.81	0.81	0.81
Coarse mass	2.94	3.82	11.40	24.50



Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.

Table 17—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity at Snoqualmie Pass, Washington

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	<i>Percent</i>	<i>Miles</i>	<i>Kilometers</i>
<4 (Baseline)	50	95.7	154.0
	60	95.7	154.0
19 (Good)	50	24.5	39.5
	60	23.6	38.0
	70	22.6	36.3
	80	21.4	34.4
	90	19.7	31.7
114 (Unhealthy for Sensitive Groups)	50	4.3	7.0
	60	4.2	6.8
	70	4.1	6.6
	80	4.0	6.5
	90	3.9	6.3
245 (Unhealthy)	50	2.0	3.2
	60	2.0	3.2
	70 to 90	1.9	3.1

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Snoqualmie Pass (Washington)





Southern Region (National Forest System—Region 8): Great Smoky Mountains National Park (Tennessee)

Particulate data from 935 days (March 1988 to May 1999) at Great Smoky Mountains National Park were chosen to represent baseline and elevated regional air quality concentrations (table 18). Table 19 shows the simulated visual range at different levels of $PM_{2.5}$ concentration (<11, 19, 114, and 245 $\mu\text{g}/\text{m}^3$) and relative humidity for Great Smoky Mountains National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<11 $\mu\text{g}/\text{m}^3$ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Great Smoky Mountains National Park (EPA 2014).

Table 18—Constituents of particulate matter ($PM_{2.5}$ and PM_{10}) at baseline (<11 $\mu\text{g}/\text{m}^3$) and elevated (19, 114, and 245 $\mu\text{g}/\text{m}^3$) levels in Great Smoky Mountains National Park, Tennessee

Particulate matter constituents	Particulate matter concentration			
	<11 (Baseline)	19	114	245
		$\mu\text{g}/\text{m}^3$		
Ammonium sulfate	6.42	13.97	13.97	13.97
Ammonium nitrate	0.43	0.31	0.31	0.31
Organic carbon	2.78	3.46	92.66	215.65
Black carbon	0.47	0.23	6.03	14.04
Fine soil	0.55	1.03	1.03	1.03
Coarse mass	5.74	7.23	11.40	24.50



Note: PM_{10} and $PM_{2.5}$ are composed of particles that are $\leq 10 \mu\text{m}$ and $\leq 2.5 \mu\text{m}$ in diameter, respectively.



Table 19—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Great Smoky Mountains National Park, Tennessee

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	Percent	Miles	Kilometers
<11 (Baseline)	60	38.0	61.2
	70	23.2	37.4
	80	19.6	31.5
19 (Good)	60	16.2	26.0
	70	4.2	6.8
	80	4.0	6.5
114 (Unhealthy for Sensitive Groups)	60	3.8	6.1
	70	2.0	3.2
	80	1.9	3.1
245 (Unhealthy)	60	1.8	2.9
	70		
	80		

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Great Smoky Mountains National Park (Tennessee)





Southern Region (National Forest System—Region 8): Mammoth Cave National Park (Kentucky)

Particulate data from 1,067 days of sampling (October 1991 to August 2003) at Mammoth Cave National Park were chosen to represent baseline and elevated regional air quality concentrations (table 20). Table 21 shows the simulated visual range at different levels of PM_{2.5} concentration (<12, 19, 114, and 245 µg/m³) and relative humidity for Mammoth Cave National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<12 µg/m³ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Mammoth Cave National Park (EPA 2014).

Table 20—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<12 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Mammoth Cave National Park, Kentucky

Particulate matter constituents	Particulate matter concentration			
	<12 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	6.94	13.99	13.99	13.99
Ammonium nitrate	0.90	0.65	0.65	0.65
Organic carbon	2.82	3.03	92.22	215.22
Black carbon	0.48	0.20	6.01	14.01
Fine soil	0.58	1.13	1.13	1.13
Coarse mass	4.43	6.26	11.40	24.50


Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.



Table 21—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Mammoth Cave National Park, Kentucky

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	Percent	Miles	Kilometers
<12 (Baseline)	50	40.3	64.8
	70	31.6	50.8
19 (Good)	50	27.5	44.2
	60	23.4	37.7
	70	19.6	31.6
	80	16.2	26.0
114 (Unhealthy for Sensitive Groups)	50	4.4	7.1
	60	4.2	6.8
	70	4.0	6.5
	80	3.8	6.1
245 (Unhealthy)	50	2.0	3.3
	60	2.0	3.2
	70	1.9	3.1
	80	1.8	2.9

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Mammoth Cave National Park (Kentucky)





Southern Region (National Forest System—Region 8): Big Bend National Park (Texas)

Particulate data from 973 days of sampling (March 1988 to May 1999) at Big Bend National Park were chosen to represent baseline and elevated regional air quality concentrations (table 22). Table 23 shows the simulated visual range at different levels of PM_{2.5} concentration (<6, 19, 114, and 245 µg/m³) and relative humidity for Big Bend National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<6 µg/m³ fine and coarse particulates) and three levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Big Bend National Park (EPA 2014).

Table 22—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<6 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Big Bend National Park, Texas

Particulate matter constituents	Particulate matter concentration			
	<6 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	2.47	4.31	4.31	4.31
Ammonium nitrate	0.24	0.42	0.42	0.42
Organic carbon	1.30	10.89	100.08	223.07
Black carbon	0.21	0.71	6.52	14.53
Fine soil	1.20	2.67	2.67	2.67
Coarse mass	7.69	11.82	11.82	24.50

Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.

Table 23—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Big Bend National Park, Texas

PM _{2.5} concentration ^a	Relative humidity		Visual range	
	Percent	Miles	Kilometers	
<6 (Baseline)	20	77.7	125.0	
	20	28.0	45.1	
19 (Good)	30	27.7	44.5	
	40	27.0	43.5	
	50	25.7	41.3	
	60	24.2	38.9	
	70	22.5	36.2	
	70	22.5	36.2	
114 (Unhealthy for Sensitive Groups)	20	4.7	7.5	
	30	4.6	7.4	
	40	4.5	7.2	
	50	4.4	7.1	
	60	4.3	6.9	
	70	4.2	6.7	
	70	4.2	6.7	
245 (Unhealthy)	20	2.2	3.5	
	30 to 40	2.1	3.4	
	50 to 60	2.0	3.2	
	70	1.9	3.1	

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Big Bend National Park (Texas)





Eastern Region (National Forest System—Region 9): Acadia National Park (Maine)

Particulate data from 986 days of sampling (March 1988 to May 1999) at Acadia National Park were chosen to represent baseline and elevated regional air quality concentrations (table 24). Table 25 shows the simulated visual range at different levels of PM_{2.5} concentration (<6, 19, 114, and 245 µg/m³) and relative humidity for Acadia National Park. The simulated images show a baseline representing an area free from smoke-impaired visibility (<6 µg/m³ fine and coarse particulates) and two levels of impairment (table 1). Data used for estimating the effect of relative humidity on visual range during the May to September fire season are from Acadia National Park (EPA 2014).

Table 24—Constituents of particulate matter (PM_{2.5} and PM₁₀) at baseline (<6 µg/m³) and elevated (19, 114, and 245 µg/m³) levels in Acadia National Park, Maine

Particulate matter constituents	Particulate matter concentration			
	<6 (Baseline)	19	114	245
		µg/m ³		
Ammonium sulfate	3.07	6.83	6.83	6.83
Ammonium nitrate	0.37	0.71	0.71	0.71
Organic carbon	1.59	10.42	99.61	222.60
Black carbon	0.34	0.68	6.49	14.50
Fine soil	0.22	0.36	0.36	0.36
Coarse mass	4.66	5.78	11.40	24.50


Note: PM₁₀ and PM_{2.5} are composed of particles that are ≤10 µm and ≤2.5 µm in diameter, respectively.


Table 25—Simulated visual range as a function of PM_{2.5} concentration (µg/m³) and relative humidity in Acadia National Park, Maine

PM _{2.5} concentration ^a	Relative humidity	Visual range	
	Percent	Miles	Kilometers
<6 (Baseline)	50	66.5	107.0
	70	20.7	33.3
19 (Good)	50	25.0	40.2
	60	22.9	36.9
	70	20.7	33.3
	80	18.4	29.6
114 (Unhealthy for Sensitive Groups)	50	4.3	7.0
	60	4.2	6.8
	70	4.1	6.5
	80	3.9	6.3
245 (Unhealthy)	50 to 60	1.9	3.1
	70	1.8	2.9
	80	1.7	2.7

^a PM_{2.5} is composed of particles that are ≤2.5 µm in diameter.

Acadia National Park (Maine)





Pacific Northwest Research Station

Web site	http://www.fs.fed.us/pnw
Telephone	(503) 808-2592
Publication requests	(503) 808-2138
FAX	(503) 808-2130
E-mail	pnw_pnwpubs@fs.fed.us
Mailing address	Publications Distribution Pacific Northwest Research Station P.O. Box 3890 Portland, OR 97208-3890



Federal Recycling Program
Printed on Recycled Paper

U.S. Department of Agriculture
Pacific Northwest Research Station
1220 SW 3rd Ave., Suite 1400
P.O. Box 3890
Portland, OR 97208-3890

Official Business
Penalty for Private Use, \$300